



## EXPLORING THE SOIL-ASSOCIATED BACTERIAL MICROBIOME OF COFFEE PLANTATIONS IN DIFFERENT REGIONS OF COLOMBIA: A METABARCODING APPROACH †

[EXPLORACIÓN DEL MICROBIOMA BACTERIANO ASOCIADO AL SUELO DE PLANTACIONES DE CAFÉ EN DIFERENTES REGIONES DE COLOMBIA: UN ENFOQUE DE METABARCODING]

Lorena Jacqueline Gómez-Godínez<sup>1</sup>, Víctor Ochoa<sup>2</sup>, Valeria Faggioli<sup>3</sup> and Marco Cristancho<sup>2\*</sup>

<sup>1</sup>Laboratorio de Recursos Genéticos Microbianos, Centro Nacional de Recursos Genéticos. Instituto Nacional de Investigación Forestales, Agrícolas y Pecuarias. Boulevard de la Biodiversidad 400, Rancho las Cruces, C.P. 47600. Tepatitlán de Morelos, Jalisco, México

<sup>2</sup>Plant Pathology Department, National Center for Coffee Research- CENICAFE, Pedro Uribe Mejía, km 4 vía Chinchiná-Manizales, C.P. 170009. Chinchiná-Manizales, Colombia. Email: [marco.cristancho@cafedecolombia.com](mailto:marco.cristancho@cafedecolombia.com)

<sup>3</sup>Instituto Nacional de Tecnología Agropecuaria INTA, Av. Rivadavia 1439. Ciudad Autónoma de Buenos Aires. C.P. C1033AAE. Buenos Aires, Argentina

\*Corresponding author

### SUMMARY

**Introduction:** Coffee is one of the leading tropical crops produced worldwide. Colombia ranks third in coffee production in the world. Microorganisms associated with coffee plants can have many biotechnological applications, such as plant growth promotion and biological control. **Objective:** To describe the bacteria present in the soil associated with coffee cultivation. This was done using a bulk sequencing or 16S rRNA metabarcoding approach. **Methodology:** Soil samples were collected to analyze their microbiome from three different departments of Colombia (Cauca, Risaralda and Magdalena). The plants were of different ages, and the crop management was different (conventional and organic). Subsequently, an analysis was carried out using Qiime2 to describe the communities associated with coffee cultivation and soil chemical properties. **Results:** Some important genera were identified, such as *Janthinobacterium*, *Bacillus*, *Actinomadura* and *Actinoallomurus*. These genera can be used as plant growth promoters, organic matter transformers, producers of antibiotics and metabolites with potential biotechnological applications. **Implications:** The study presented in this manuscript describes the communities associated with the soil in coffee cultivation. However, a much more complex approach could be through metagenomics, where the bacterial communities and the functions of; however, this vision is up to ten times higher in cost. **Conclusion:** Through the results obtained, it is concluded that the studies carried out through metabarcoding help to understand the composition of the microorganisms associated with coffee and, with this, try to elucidate the functions of the associated microorganisms.

**Key words:** Metabarcoding; coffee; uncultured bacteria species.

### RESUMEN

**Introducción:** El café es uno de los principales cultivos tropicales producidos a nivel mundial. Colombia ocupa el tercer lugar en producción de café en el mundo. Los microorganismos asociados a las plantas de café pueden tener una gran cantidad de aplicaciones biotecnológicas, como la promoción de crecimiento vegetal y el control biológico. **Objetivo:** Describir las bacterias presentes en el suelo asociadas al cultivo del café. Esto se hizo mediante un enfoque de secuenciación masiva o metabarcoding del gen ARNr 16s. **Metodología:** Se recolectaron muestras de microbioma del suelo de tres departamentos diferentes de Colombia (Cauca, Risaralda y Magdalena). Las plantas tuvieron diferentes edades y el manejo del cultivo fue diferente (convencional y orgánico). Posteriormente se realizó un análisis mediante Qiime2 para describir las comunidades asociadas al cultivo de café. **Resultados:** Se identificaron algunos Phylum importantes como, por ejemplo, Acidobacterias, Actinobacterias, Firmicutes y Verrucomicrobia, Dentro de

† Submitted September 28, 2023 – Accepted February 27, 2024. <http://doi.org/10.56369/tsaes.5196>



Copyright © the authors. Work licensed under a CC-BY 4.0 License. <https://creativecommons.org/licenses/by/4.0/>

ISSN: 1870-0462.

ORCID = Lorena Jacqueline Gómez-Godínez: <http://orcid.org/0000-0002-9759-7207>, Valeria Faggioli: <http://orcid.org/0000-0002-7066-6866>, Marco Cristancho: <http://orcid.org/0000-0001-9115-4117>

los géneros importantes, se lograron identificar a *Janthinobacterium*, *Bacillus*, *Actinomadura* y *Actinoallomurus*. Estos géneros pueden ser utilizados como promotores del crecimiento vegetal, transformadores de materia orgánica, productores de antibióticos y metabolitos con potenciales aplicaciones biotecnológicas. **Implicaciones:** El estudio presentado en este manuscrito es descriptivo, de las comunidades asociadas al suelo en el cultivo de café, sin embargo, un enfoque mucho más complejo podría ser a través de metagenómica donde se puede describir a las comunidades de bacterias y las funciones de estas, sin embargo, esta visión es hasta diez veces más elevada en cuestiones de costo. **Conclusión:** A través de los resultados se concluye que los estudios realizados a través de metabarcoding ayudan a entender la composición de los microorganismos asociados al café y con esto tratar de dilucidar las funciones de los microorganismos asociados.

**Palabras clave:** metabarcoding; café; especies de bacterias no cultivadas.

## INTRODUCTION

Coffee is the second most consumed beverage after water and the most traded tropical agricultural product in the world (Mussatto *et al.*, 2011); it is a perennial plant belonging to the Rubiaceae family. The *Coffea* genus consists of about 103 species, where the *Coffea arabica* species is the most used for the production of beverages worldwide (Davis *et al.*, 2006). Colombia is one of the world's leading coffee producers, ranking third after Brazil and Vietnam. Within Colombia, Cauca is the fourth-largest coffee-producing department, Risaralda is sixth, and Magdalena is fourteenth. Soils are one of the most dynamic ecosystems on the planet. They are home to a diverse community of microorganisms, including bacteria, fungi, viruses, archaea, and protozoa. These microorganisms interact with each other and with the soil environment to perform essential ecosystem functions (Lladó *et al.*, 2017).

The study of plant-associated microorganisms is important for a number of biotechnological applications, such as the biological control of plant pathogens, the promotion of plant growth, and the isolation of active compounds. Biodiversity characterization has also been used to monitor and assess the state of the environment (Bonatelli *et al.*, 2021; Lahlali *et al.*, 2022; Lopes *et al.*, 2021). Changes in the structure of the microbial community can influence the stability of populations in the face of disturbances. Therefore, the identification and conservation of soil microbial components is essential for resilience under changing conditions, as well as to support current and future food security (Bertola *et al.*, 2021).

The importance of identifying the microorganisms associated with coffee is a crucial step to allow the prospecting of beneficial microorganisms that can be used as biofertilizers in soils and starter cultures to improve the quality of the coffee drink since it is known that the profile microbial promotes changes in the sensory and chemical profile in coffee fermentation (Junior *et al.*, 2021; Duong *et al.*, 2020). Furthermore, coffee cultivars are more adapted to some regions than others, and their associated microorganisms may also be shaped by climatic and edaphic characteristics,

which may be related to the sensory experience obtained from this product (terroir) (Veloso *et al.*, 2020; Williams *et al.*, 2022).

One of the factors that can modify the communities of microorganisms in the soil is tillage. It is known that conventional tillage causes physical alterations of the upper horizon of the soil and can decrease organic matter (Brangarí *et al.*, 2022). On the other hand, zero tillage and organic management are practices that can favor the enrichment of soil with organic matter and nutrients, reduce soil erosion, increase water availability, and improve the stability of the soil system in the face of changes in humidity and temperature (Sun *et al.*, 2018; Schmidt *et al.*, 2018; Xin *et al.*, 2018). It has been shown that various agronomic management can significantly alter soil biomass, as well as the composition of the microbial communities present. For example, greater microbial diversity and more microbial biomass have been reported in soil treated under no-till management compared to soils with conventional management (Wang *et al.*, 2017; Yin *et al.*, 2017; Mathew *et al.*, 2012; Guo *et al.*, 2016).

Most studies on crop-associated bacteria and their community structure have been conducted using culture-dependent approaches. However, since a high percentage of natural bacteria remain in a non-culturable state (Bodor *et al.*, 2020), the use of culture-independent methods provides additional information on the diversity of bacterial communities (Rincon-Florez *et al.*, 2013).

In the last decade, the use of high-throughput sequencing has substantially improved our understanding of the global biogeography and ecology of soil bacterial and fungal communities (Howe *et al.*, 2021). This strategy, along with metabarcoding, represents a fast, accurate, and cost-effective technology for testing thousands of environmental samples simultaneously.

The objective of this study was to analyze the communities associated with coffee, of different ages and grown in different regions of Colombia, as well as under different agricultural systems, conventional and organic, using a crop-independent approach.

## MATERIAL AND METHODS

### Description of the study site


The study was conducted in three Colombian coffee-growing regions: El Tambo (Cauca), Quinchía (Risaralda), and Minca (Magdalena). A total of 20 coffee farms were sampled, 8 in Cauca, 8 in Risaralda, and 4 in Magdalena. The soil samples were collected between July and August 2021, under the climatic and environmental conditions of each region. Around the coffee-plant, the surface of each area was cleaned to remove any organic material present. A sterile metal spatula was used to collect approximately 50-150 g of soil (a mixture of soil, sand, and small soil grains) from a depth of 0.2 meters, the distance between coffee plant for sampling was 1.5 meters. Five different coffee plants samples were collected from each farm, for a total of 30 soil samples. The samples were stored in sterile Ziploc-type plastic bags and transported to the Laboratory of Molecular Interactions of Microorganisms in Agriculture - LIMMA, located at the Universidad de Los Andes, for further processing (Figure 1).

### Physical-chemical soil analysis

The physicochemical analyses were performed on 100 g of each sample at the Soil and Foliar Laboratory of the Technological University of Pereira, Colombia.

The pH was measured in water (1:1 w/v aqueous extract) using a potentiometer. The Organic Matter (OM) was determined with the Walkley-Black photometric method. Electrical Conductivity (EC) was measured in paste saturated with water using a conductivity meter (Keren 2016; Kalra and Yash 1995). The following ions were determined using the atomic absorption technique (Acetate+EDTA): Iron (Fe), Manganese (Mn), Zinc (Zn), and Copper (Cu). Boron (B) was determined by extraction with monocalcium phosphate, azomethine, and photometric detection (Maurya *et al.*, 2018). Sulfur (S) was determined by the turbidimetric method under extraction with monocalcium phosphate (Wall *et al.*, 1980). Phosphorus (P) was analyzed with the Bray II photometric method. Aluminum was also measured (Wuenschel *et al.*, 2015; Lin and Coleman *et al.*, 1960).

The texture was determined by Bouyoucos and the exchangeable acidity by the volumetric method under extraction with KCl (Beretta *et al.*, 2014). The bases Potassium (K), Calcium (Ca), Magnesium (Mg), and Sodium (Na) were analyzed by the ammonium acetate method, using the atomic absorption technique. Solubles were measured in water-saturated paste, using the same technique. Finally, the Cation Exchange Capacity (CEC) was analyzed volumetrically (Sumner *et al.*, 1996; Soylak *et al.*, 2003; Yunan *et al.*, 2028).



**Figure 1.** Graphical overview of the methods of this study.

## Sample processing and metagenomic DNA extraction

Each Ziploc bag was tempered before processing. Using sterile metallic mini-spatulas, 250 mg of soil was taken and added to a sterile 2 ml Eppendorf tube. DNA was extracted using the DNeasy® PowerSoil® Pro-Kit (Qiagen) and EZ-10 Spin Column Soil DNA Miniprep Kit (BioBasic) following the manufacturer's instructions. Possible contaminants were removed with the kit's own resins, and then the DNA was separated with silica filters. The DNA extraction was performed individually for each replicate. DNA was quantified using spectrophotometry with the NanoDrop™ 2000 (Thermo Fisher Scientific™) and Qubit 2.0 (Thermo Fisher Scientific™) kits following the manufacturer's protocols. The integrity of the DNA was observed by electrophoresis in a 1% agarose gel. There were 6 conditions with 3 or 5 replicates per condition, having a total of 25 samples for metagenomic sequencing. These were stored at -80°C in sterile Eppendorf tubes until further processing (Figure 1).

## Sequencing of the metagenome

Libraries were constructed using the Illumina Metagenomics Sequencing technique and the MiSeq PE250 (Matched Ends 2x250) Sequencing Kit. The 16S V3/V4 region of bacteria was amplified using primers 341F (5'-CCTAYGGGRBGCASCAG-3') and 806R (5'-GGACTACNNGGGTATCTAAT-3') (Yu *et al.*, 2005). All sequencing was performed by external services of the National Institute of Agricultural Technology - Marcos Juárez Agricultural Experimental Station, in Argentina (Figure 1).

## Bioinformatics Analysis

Sequence quality was assessed using FastQC (v0.11.9) (Andrews, 2010). Quality control and processing were performed using QIIME2 (v2021.4.0) (Bolyen *et al.*, 2019). Sequences were demultiplexed using a Q-score of 30, Illumina adapters, and poor sequencing quality were removed with cutadapt and DADA2 (Callahan *et al.*, 2016). Clean sequences were compared to the greengenes database (gg-13-8-99-515-806-nb-classifier.qza), and taxonomic assignment (ASVs) was performed. Alpha diversity indices (observed species, Chao1, Shannon, and Simpson) were calculated from the rarefaction curves. Beta diversity indices were calculated using Euclidean distance and UniFrac (weighted and unweighted) and visualized through principal coordinate analysis (PCoA) (Figure 1).

## RESULTS

### Soil analysis

Soils from three colombian regions (Cauca, Risaralda, and Magdalena) were characterized. The pH of the soils varied, with Cauca having the lowest pH of 4.5, followed by Risaralda 4.9 and 5.0 to Magdalena (Table 1). The percentage of organic matter also varied, with Risaralda\_9 having the lowest percentage (7.9%), followed by Risaralda\_15 (8.7%), Risaralda\_12 (14.6%), and Cauca (23%). The nitrogen content varied, with Risaralda\_9 having the lowest percentage (0.33%), followed by Risaralda\_15 (0.42%), Risaralda\_12 (0.58%), and Cauca (0.72%). The percentage of phosphorus was highest in Risaralda\_9 (32%), followed by the other samples (4%). The percentage of potassium was similar across all samples, ranging from 1% to 1.5%. The site with the highest amount of iron was Risaralda 12, with 218 parts per million (ppm), followed by Cauca with 169 ppm; Manganese oscillated between 31 ppm (Risaralda\_12), and the lowest concentration was presented in Cauca and Risaralda\_15 with 18 ppm. Zinc was found in a concentration of 14 ppm in Cauca and 4 ppm in Risaralda\_15 (Table 1). The samples from this site are divided into Magdalena\_17 (30-year-old plants, conventional management) and Magdalena\_19 (new plants, conventional management). Samples from Magdalena\_19 were not analyzed for physical-chemical properties; this is denoted as NA in Table 1. The main reason the analysis was not carried out was due to the conditions of the sample (soil texture characteristics, made DNA extraction difficult) and the fact that there was no opportunity to repeat the sampling due to difficult access to the place.

### Profile of bacterial communities associated with coffee cultivation

Amplicons from 25 samples corresponding to the different treatments were obtained and analyzed as described above. A total of 2,649,699 reads were obtained, with an average of 115,204 reads per sample. After filtering, 1,901,635 reads were obtained and 132 ASVs were identified. These were assigned to 14 phyla, 24 classes, 14 orders, 23 families, and 42 bacterial genera. The diversity indices, calculated from the rarefaction curves, remained constant across the different treatments.

The most abundant phylum in the soil samples from the Cauca department was Firmicutes. Chloroflexi and Acidobacteria were the next most abundant phyla. In the soil samples from the Risaralda department, Acidobacteria was the most abundant phylum, followed by Nitrospirae and Proteobacteria. In the soil samples from the Magdalena department, the most

abundant phyla were Planctomycetes, Acidobacteria, and Verrucomicrobia. The results are summarized in Figure 2.

In most samples, the most abundant family was Hyphomicrobiaceae, however, in the samples from Risaralda\_19, and Risaralda\_9, the most abundant family was Koribacteraceae, and Pseudomonadaceae, respectively. The Rhodospirillaceae family was the second most abundant in the samples from Cauca, Magdalena\_17 and Magdalena\_19. In the case of the samples from Risaralda\_9 and Risaralda\_12, the second most abundant family was Micrococcaceae, and in the case of the sample from Risaralda\_15 Koribacteraceae. Koribacteraceae was the third family in the samples from Cauca and Magdalena\_17, in the samples from Risaralda\_15 and Magdalena\_19 the family Syntrophobacteraceae and Sinobacteraceae, ranked third respectively (Figure 3).

Finally, some of the genera identified in Cauca were *Conexibacter*, *Bacillus*, *Actinoallomurus*, *Rhodomicrobium*, *Actinomadura*, and *Burkholderia*. In the Risaralda\_9 samples, *Chryseobacterium*, *Pedobacter*, *Janthinobacterium*, *Pseudomonas* and *Rhodococcus* were found, in Risaralda\_12, *Arthrobacter*, *Labrys*, *Pilimelia*, *Clostridium*, and *Rhodomicrobium*, in Risaralda\_15, *Anaeromyxobacter*, *Steroidobacter*, *Nitrospira*, *Mesorhizobium* and *Pedomicrobium*. In Magdalena\_17, *Nostocoida*, *Conexibacter*, *Actinomadura*, *Burkholderia* and *Edaphobacter*, and

finally in Magdalena\_19 *Novosphingobium*, *Kaistobacter*, *Methylibium*, *Flavobacterium* and *Phenylobacterium* (Figure 4).


The analysis of beta diversity, based on PCoA, did not allow us to corroborate differences in the structure of the microbiota associated with the coffee soil in the different regions of Colombia (Figure 5).

## DISCUSSION

Soils intended for agricultural crops are often low in organic matter, with contents of less than 4% to 6% (Hatten & Liles 2019). However, this is not the case for coffee crops, which are typically grown under shade and benefit from the mulching effect of leaf litter, finding percentages of organic matter from 4% to 11% (López-Báez *et al.*, 2016; Darfīs and Putri 2021). This helps to protect the soil from erosion and enhances the accumulation of organic matter. In our study, we found that the organic matter content of coffee soils ranged from 5.5% to 23%. Organic matter is important for plant growth because it provides nutrients, such as nitrogen (Zheng *et al.*, 2023). In Colombia, mathematical models have been developed to estimate the nitrogen content of coffee soils based on the organic matter content (Salazar *et al.*, 2023). Our results confirm this relationship, as we found that the soil samples from Cauca, which had the highest organic matter content (23%), also had the highest nitrogen content (0.72%).

**Table 1. Description of study sites and soil chemical properties. Where: pH: potential of hydrogen, N: Nitrogen, OM: Organic Matter, CEC: Cation exchange capacity, K: Potassium, Ca: Calcium, Mg: Magnesium, Al: Aluminum, Fe: Iron, Mn: Manganese, Zn: Zinc, Cu: Copper, B: Boron, S: Sulfur, P: Phosphorus and NA: Does not apply.**


ID and region	Age of plant	Management	Latitude	Longitude	pH	% N total	% OM	CEC (meq/100g)	K (meq/100g)	Ca <sup>2+</sup> (meq/100g)	Mg <sup>2+</sup> (meq/100g)	Al <sup>3+</sup> (meq/100g)	Fe <sup>2+</sup> (ppm)	Mn <sup>2+</sup> (ppm)	Zn <sup>2+</sup> (ppm)	Cu <sup>2+</sup> (ppm)	B (ppm)	S <sup>2-</sup> (ppm)	P (ppm)
Cauca	<1	organic	2.461	-76.8	4.5	0.72	23	51	1.4	2	1	4	169	18	14	3	0.1	11	4
Risaralda_9	10	organic	5.352	-75.73	5.8	0.33	7.9	48	1.5	13	3.3	NA	66	25	8	2	0.2	11	32
Risaralda_12	5	organic	5.349	-75.73	4.9	0.54	15	48	1	6	2.3	1.2	72	31	9	3	0.1	12	4
Risaralda_15	5	organic	5.364	-75.73	5	0.36	8.7	34	1.3	8	1.9	1.2	218	18	4	4	0.4	10	9
Magdalena_17	30	conventional	11.077	-74.04	5	0.25	5.5	NA	0.1	1.1	0.5	2.3	NA	NA	NA	NA	NA	NA	2
Magdalena_19	<1	conventional	11.088	-74.08	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA



**Figure 2.** Phylum bacterial communities associated with soil samples obtained from different regions of Colombia. The bars represent the relative abundance. Sample 1 a,c,d: samples obtained from the department of Cauca in the municipality of Tambo, 4-month-old plants with organic management; sample 9 a,b,d: samples obtained from the department of Risaralda in the municipality of Quinchia, 10-year-old plants with organic management; sample 12 a,b,c,d and e: samples obtained from the department of Risaralda in the municipality of Quinchia, 5-year-old plants with organic management; sample 15 a,c,d and e: samples obtained from the department of Risaralda in the municipality of Quinchia, 5-year-old plants with organic management; sample 17 a,b,c,d and e: samples obtained from the department of Magdalena in the municipality of Minca, 30-year-old plants with conventional management; sample 19 a,b,c,d and e: samples obtained from the department of Magdalena in the municipality of Minca, new plants with organic management.

Microorganisms are the living component of soil, and their abundance and activity depend on a variety of factors, such as soil type and management, plant type and growth, and root exudates. In this study, we report the diversity of bacteria associated with coffee plants of different ages, under conventional and organic management. We found that the Acidobacteria phylum was abundant in samples from the Cauca

department, Risaralda\_9, Magdalena\_17, and Magdalena\_19. The Acidobacteria phylum is positively correlated with organic matter and is involved in carbon sequestration. Additionally, Acidobacteria dominate soils with high organic matter content and are involved in the degradation of lignocellulose (Rawat *et al.*, 2012).




**Figure 3.** Family bacterial communities associated with soil samples obtained from different regions of Colombia. The bars represent the relative abundance.

On the other hand, the Firmicutes phylum was most abundant in the soils of Cauca in comparison with Risaralda and Magdalena sites. This phylum is typically found in soil, but in lower abundance than other phyla (Youssef and Elshahed, 2008). Firmicutes include gram-positive bacteria that can form endospores, which allow them to survive in adverse environmental conditions. *Bacillus* is a genus of Firmicutes that is commonly found in soil.

In the Risaralda\_9 sample, we found a high percentage of the Proteobacteria phylum, which is common in agricultural ecosystems and plays a variety of roles in the soil (Valinsky *et al.*, 2002), including carbon, nitrogen, and sulfur cycling (Spain *et al.*, 2009). The Nitrospirae phylum was also identified in the Risaralda\_9 and Magdalena\_19 samples. This phylum is characterized by a single class, Nitrospira, and its members are aerobic chemolithotrophs that can

perform a variety of metabolic functions, including nitrification, sulfate reduction, and magnetotaxis. Some species of Nitrospira are important participants in the nitrogen cycle (Hu., *et al.*, 2021).

Another interesting and consistent phylum found in the samples was Verrucomicrobia. It was found in high percentages in the samples from Risaralda\_12, Magdalena\_17, and Magdalena\_19. Verrucomicrobia are difficult to cultivate, and most are facultative mesophiles or strict anaerobes (Bergmann *et al.*, 2011). They are saccharolytic and oligotrophic. Little is known about their function, but some representatives of Verrucomicrobia associated with rice plants have been found to incorporate plant-derived carbon for growth (Hernández *et al.*, 2015). This may explain the high abundance of Verrucomicrobia in the Risaralda\_12 samples, which had the highest percentage of organic matter.




**Figure 4.** Family bacterial communities associated with soil samples obtained from different regions of Colombia. The bars represent the relative abundance.

The phylum Chlamydiae is a small group of obligate intracellular bacteria that were once thought to only infect humans and other animals (Elwell *et al.*, 2016). However, recent research has shown that Chlamydiae are actually much more widespread and can be found in a wide variety of host organisms, including protists, arthropods, and vertebrates. Chlamydiae have also been found in a variety of environmental samples, suggesting that they may play a role in the cycling of nutrients in the environment.

The most abundant genera in the samples from Cauca were *Conexibacter*, *Bacillus*, and *Actinoallomurus*. *Conexibacter* is a genus of Actinobacteria that is typically found in acidic soils, such as those in the Cauca region (pH 4.5). These bacteria are known to contribute to the carbon cycle in soil environments and

may also play a role in the nitrogen cycle (Pukall *et al.*, 2010; Seki *et al.*, 2012). *Bacillus* is another genus that was found in the samples. This genus is common in coffee plants and some species have been shown to inhibit the growth of pathogenic fungi, such as *Colletotrichum gloeosporioides* and *Fusarium oxysporum* (Kejela *et al.*, 2016). *Bacillus* species also have a variety of other beneficial properties, including the ability to promote plant growth, solubilize and mineralize nutrients, produce phytohormones, and secrete antimicrobial compounds (Saxena *et al.*, 2020). *Actinoallomurus* is the third most abundant genus in the samples from Cauca. These bacteria are of great interest due to their secondary metabolism, which could be used to find organisms with potential in the production of antibiotics against resistant pathogens (Pozzi *et al.*, 2010).



**Figure 5.** Calculated beta diversity for bacterial communities associated with coffee soil samples.

In the samples from Risaralda\_9 we found in greater abundance the genera *Chryseobacterium*, *Pedobacter*, *Janthinobacterium*. *Chryseobacterium* has been reported in different environments such as wastewater, clinical samples, and soil (Kämpfer *et al.*, 2003, 2009; Shen *et al.*, 2005). The strains that represent this genus are Gram-negative, immobile, and yellow in color. Some species have been reported as growth promoting bacteria (Chhetri *et al.*, 2022). Bacteria within the *Pedobacter* genus are characterized by being resistant to antibiotics (Bjerketorp *et al.*, 2021), and some species within this genus have been characterized by breaking down cellulose (Zhou *et al.*, 2019). Within *Janthinobacterium* we find gram-negative bacteria, producers of some substances such as violacein and capable of inhibiting some pathogenic fungi such as *Pythium ultimum*, *Rhizoctonia solani* and *Rhizoctonia oryzae*, in crops such as wheat (Yin *et al.*, 2021). It has been reported that when a soil is enriched with bacteria of this genus, it shows the ability to be less susceptible to dollar spot caused by the pathogenic fungus *Clarireedia spp.* (Chou *et al.*, 2021). Finally, *Janthinobacterium* has been shown to promote plant growth, thanks to the production of indole-3-acetic acid (Kuffner *et al.*, 2008).

The most abundant genera in the Risaralda\_12 samples are *Arthrobacter*, *Labrys*, and *Pilimelia*. *Arthrobacter* is a genus of obligate aerobic bacteria that grow in a rod-coccus cycle (Gobbetti and Rizzello, 2014). They are commonly found in soils, aerial surfaces of plants,

and sewage sediments. They do not form endospores and are highly proteolytic. Some species of *Arthrobacter*, such as *Arthrobacter agilis*, have been reported as plant growth promoting bacteria. *A. agilis* was isolated from corn and has the ability to solubilize iron and increase the biomass of beans and *Medicago sativa* (Valencia-Cantero *et al.*, 2007; Velázquez-Becerra *et al.*, 2011).

*Labrys* is a genus of aerobic, gram-negative, motile, and non-sporulating bacteria. Their shape varies from spherical to short rods. They were first described by Vasil'eva and Semenov (1984). Some species of *Labrys*, such as *Labrys metilaminiphilus*, can use methanol as a carbon source (Miller *et al.*, 2005). *Pilimelia* is a genus of bacteria within the Actinobacteria. They produce branched, septate substratum hyphae, but the aerial mycelium does not develop. All *Pilimelia* species are aerobic, mesophilic, and gram-positive (Schaefer, 1973). Some *Pilimelias* have the ability to degrade plant polymers such as starch, cellulose, hemicellulose, pectin, or lignin. This may be due to the litter present in coffee crops.

The main genera in the Risaralda\_15 samples were *Anaeromyxobacter*, *Steroidobacter*, and *Nitrospira*. *Anaeromyxobacter* are Fe (III) reducers, meaning they can reduce the oxidized form of iron (Fe(III)) to the reduced form (Fe(II)) (Sanford *et al.*, 2002; Wang *et al.*, 2020). This process is important in the cycling of iron in the environment. *Steroidobacter* are found in

soil, specifically organic soybean soils. They are believed to have a role in carbon decomposition, which is the process by which organic matter is broken down into smaller molecules (Lian *et al.*, 2017). *Nitrospira* are found in acid soils and have an important role in nitrification. Nitrification is the process by which ammonia is oxidized to nitrate (Séneca *et al.*, 2020). The abundance of each genus was affected by the environmental conditions. For example, the abundance of *Anaeromyxobacter* was favored in rice cultures enriched with ferrihydrite and goethite. The abundance of *Nitrospira* decreased due to drought.

The genera *Nostocoida*, *Conexibacter*, and *Actinomadura* were abundantly found in the Magdalena\_17 samples. *Nostocoida* are bacteria that were isolated from activated sludge (this bacterium needs nitrogen to grow). (Blackall *et al.*, 2000). *Conexibacter* are bacteria that are common in soil. They may have a role in the decomposition of organic material. *Actinomadura* are bacteria that are common in soil. They can produce antibiotics, vitamins, enzymes, and other valuable biologically active substances (Trüper *et al.*, 1992).

The Magdalena\_19 samples contained genes from the genera *Novosphingobium*, *Kaistobacter*, and *Methylibium*. *Novosphingobium* are bacteria that have been found as endophytes of some crops, such as rice. Some species have growth promotion potential, and one has been identified as diazotrophic, meaning it can fix nitrogen (Addison *et al.*, 2007; Gao *et al.*, 2015; Zhang *et al.*, 2016). *Kaistobacter* are bacteria that are favored when substances are added to the soil to protect plants against fungal diseases (Baćmaga *et al.*, 2021). *Methylibium* were a genus described by Nakatsu *et al.* (2006). Many members of this genus hydrolyze urea, and some species have also shown their ability to biodegrade some undesirable compounds in biogas production (Boada *et al.*, 2020; Nakatsu *et al.*, 2006).

We found several genera of bacteria with great biotechnological potential, including *Janthinobacterium*, *Bacillus*, *Actinomadura*, and *Actinoallomurus*. These bacteria can produce metabolites or antibiotics that be used to treat multi-resistant bacteria, promote plant growth, and inhibit pathogenic fungi (Kuffner *et al.*, 2008; Trüper *et al.*, 1992; Kejela *et al.*, 2016; Saxena *et al.*, 2020).

Numerous studies have pointed out the importance of soil-associated microbial communities in plant growth promotion. These studies have documented the capacity of bacterial genera within communities, which are responsible for producing plant growth inducers, metabolites and antibiotics, among other substances (Poria *et al.*, 2022; Katsenios *et al.*, 2022; Gómez-Godínez *et al.*, 2023). Within this work, we

postulate the ability of bacteria to colonize the roots of coffee plants and how they could establish beneficial symbiotic interactions. This symbiosis could enhance the absorption of essential nutrients by the coffee plant, thus improving the crop's ability to take advantage of nutrients efficiently, and the synthesis of compounds related to the promotion of plant growth could positively influence root development. Water absorption, resistance to adverse conditions, and the ability to inhibit pathogenic fungi could prevent soil diseases and directly affect coffee productivity. The selective suppression of pathogens the microbial communities could contribute to the overall health of the plants and, therefore, to an increase in the productivity of coffee beans (Souza *et al.*, 2015; Glick, 2012; Abdelaal *et al.*, 2021; Urgiles-Gómez *et al.*, 2021; Asad *et al.*, 2023).

Finally, it is expected that the study will not only deepen our understanding of the diversity associated with coffee plants but also lay the foundation for innovative and sustainable strategies in the controlled use of these bacterial communities, such as, using them in bioaugmentation. Moreover, the controlled introduction of beneficial bacteria can modulate the structure of the community, favoring interactions between community members, suppressing pathogens, and increasing plant growth and coffee bean production. Detailed exploration of these biotechnological potentials could have practical implications in agriculture, improving the sustainability and profitability of coffee globally.

**Funding.** This work was supported by CABANA (Capacity building for bioinformatics in Latin America) grant, on behalf of the Global Challenges Research 408 Fund [BB/P027849/1].

**Conflict of interests.** The authors declare no competing interests.

**Compliance with ethical standards.** This work does not require approval by a bioethical committee.

**Data availability.** The datasets generated during and/or analyzed during the current study are available in the European Molecular Biology Laboratory-European Bioinformatics Institute (EMBL-EBI) with the project ID PRJEB50597 (<https://www.ebi.ac.uk/>). Or by email marco.cristancho@cafedecolombia.com

**Author contribution statement (CRediT).** LJ Gómez-Godínez - Writing-original draft, Data curation and Writing-review & editing. V. Ochoa - Methodology and Writing-review & editing. V. Faggioli - Methodology and Writing-review & editing. M. Cristancho - Funding acquisition, Methodology, Project administration and Writing-review & editing.

## REFERENCES

- Abdelaal, K., AlKahtani, M., Attia, K., Hafez, Y., Király, L. and Künstler, A., 2021. The role of plant growth-promoting bacteria in alleviating the adverse effects of drought on plants. *Biology*, 10(6), pp. 520. <https://doi.org/10.3390/biology10060520>
- Addison, S.L., Foote, S.M., Reid, N.M., and Lloyd-Jones, G., 2007. *Novosphingobium nitrogenifigens* sp. nov., a polyhydroxyalkanoate-accumulating diazotroph isolated from a New Zealand pulp and paper wastewater. *International Journal of Systematic and Evolutionary Microbiology*, 57(11), pp. 2467–2471. <https://doi.org/10.1099/IJS.0.64627-0>
- Asad, S., Priyashantha, A. K. H., Tibpromma, S., Luo, Y., Zhang, J., Fan, Z., Zhao, L., Shem, K., Miu, C., Lu, L., Promputtha, I. and Karunaratna, S. C., 2023. Coffee-associated endophytes: Plant growth promotion and crop protection. *Biology*, 12(7), pp. 911. <https://doi.org/10.3390/biology12070911>
- Bácmaga, M., Wyszowska, J. and Kucharski, J., 2021. Bacterial diversity and enzymatic activity in a soil recently treated with tebuconazole. *Ecological Indicators*, 123, pp. 107373. <https://doi.org/10.1016/J.ECOLIND.2021.107373>
- Beretta, A. N., Silbermann, A. V., Paladino, L., Torres, D., Bassahun, D., Musselli, R. and García-Lamohte, A., 2014. Análisis de textura del suelo con hidrómetro: Modificaciones al método de Bouyoucus. *Ciencia e investigación agraria*, 41(2), pp. 263-271. <http://dx.doi.org/10.4067/S0718-16202014000200013>
- Bergmann, G.T., Bates, S. T., Eilers, K. G., Lauber, C. L., Caporaso, J. G., Walters, W. A., Knight, R. and Fierer, N., 2011. The under-recognized dominance of *Verrucomicrobia* in soil bacterial communities. *Soil Biology and Biochemistry*, 43(7), pp. 1450. <https://doi.org/10.1016/J.SOILBIO.2011.03.012>
- Bertola, M., Ferrarini, A., and Visioli, G., 2021. Improvement of Soil Microbial Diversity through Sustainable Agricultural Practices and Its Evaluation by -Omics Approaches: A Perspective for the Environment, Food Quality and Human Safety. *Microorganisms*, 9, pp. 1400. <https://doi.org/10.3390/MICROORGANISM9071400>
- Bjerketorp, J., Levenfors, J. J., Nord, C., Guss, B., Öberg, B. and Broberg, A., 2021. Selective Isolation of Multidrug-Resistant *Pedobacter* spp., Producers of Novel Antibacterial Peptides. *Frontiers in Microbiology*, 12, pp. 366. <https://doi.org/10.3389/FMICB.2021.642829>
- Blackall, L.L., Seviour, E.M., Bradford, D., Rossetti, S., Tandoi, V., and Seviour, R.J., 2000. “*Candidatus Nostocoida limicola*”, a filamentous bacterium from activated sludge. *International Journal of Systematic and Evolutionary Microbiology*, 50(2), pp. 703–709. <https://doi.org/10.1099/00207713-50-2-703>
- Boada, E., Santos-Clotas, E., Bertran, S., Cabrera-Codony, A., Martín, M.J., Bañeras, L. and Gich, F., 2020. Potential use of *Methylobium* sp. as a biodegradation tool in organosilicon and volatile compounds removal for biogas upgrading. *Chemosphere*, 240, 124908. <https://doi.org/10.1016/J.CHEMOSPHERE.2019.124908>
- Bodor, A., Bounedjoum, N., Vincze, G.E., Erdeiné Kis, Á., Laczi, K., Bende, G., Szilágyi, Á., Kovács, T., Perei, K. and Rákhely, G., 2020. Challenges of unculturable bacteria: environmental perspectives. *Reviews in Environmental Science and Bio/Technology*, 19(1), pp. 1–22. <https://doi.org/10.1007/S11157-020-09522-4>
- Bonatelli, M. L., Lacerda-Júnior, G. V., dos Reis Junior, F. B., Fernandes-Júnior, P. I., Melo, I. S. and Quecine, M. C., 2021. Beneficial Plant-Associated Microorganisms From Semiarid Regions and Seasonally Dry Environments: A Review. *Frontiers in Microbiology*, 11, 3331. <https://doi.org/10.3389/FMICB.2020.553223>
- Brangarí, A. C., Lyonard, B. and Rousk, J., 2022. Soil depth and tillage can characterize the soil microbial responses to drying-rewetting. *Soil Biology and Biochemistry*, 173, p. 108806. <https://doi.org/10.1016/j.soilbio.2022.108806>
- Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P., 2016. DADA2: High resolution sample inference from Illumina amplicon data.

- Nature Methods, 13(7), 581.  
<https://doi.org/10.1038/NMETH.3869>
- Chhetri, G., Kim, I., Kim, J., So, Y. and Seo, T., 2022. *Chryseobacterium tagetis* sp. nov., a plant growth promoting bacterium with an antimicrobial activity isolated from the roots of medicinal plant (*Tagetes patula*). *The Journal of Antibiotics*, 75(6), pp. 312–320. <https://doi.org/10.1038/s41429-022-00525-7>
- Chou, M. Y., Shrestha, S., Rioux, R. and Kocha, P., 2021. Hyperlocal Variation in Soil Iron and the Rhizosphere Bacterial Community Determines Dollar Spot Development in Amenity Turfgrass. *Applied and Environmental Microbiology*, 87(10), pp. 1–20. <https://doi.org/10.1128/AEM.00149-21>
- Darfis, I. and Putri, E.L., 2021. Performance of some soil physical properties of arabica coffee plantation in solok regency. *IOP Conference Series: Earth and Environmental Science*, 741, p. 012028. <http://doi.org/10.1088/1755-1315/741/1/012028>
- Davis, A.P., Govaerts, R., Bridson, D.M., and Stoffelen, P., 2006. An annotated taxonomic conspectus of the genus *Coffea* (Rubiaceae). *Botanical Journal of the Linnean Society*, 152(4), pp. 465–512. <https://doi.org/10.1111/J.1095-8339.2006.00584.X>
- Duong, B., Marraccini, P., Maeght, J.L., Vaast, P., Lebrun, M. and Duponnois, R., 2020. Coffee microbiota and its potential use in sustainable crop management. A review. *Frontiers in Sustainable Food Systems*, 4, p. 607935. <https://doi.org/10.3389/fsufs.2020.607935>
- Elwell, C., Mirrashidi, K., and Engel, J., 2016. Chlamydia cell biology and pathogenesis. *Nature Reviews. Microbiology*, 14(6), 385. <https://doi.org/10.1038/NRMICRO.2016.30>
- Gao, S., Zhang, Y., Jiang, N., Luo, L., Li, Q. X., and Li, J., 2015. *Novosphingobium fluoreni* sp. nov., isolated from rice seeds. *International Journal of Systematic and Evolutionary Microbiology*, 65(5), pp. 1409–1414. <https://doi.org/10.1099/IJS.0.000111>
- Glick, B. R., 2012. Plant growth-promoting bacteria: mechanisms and applications. *Scientifica*, <https://doi.org/10.6064/2012/963401>
- Gobbetti, M., and Rizzello, C. G., 2014. *Arthrobacter*. *Encyclopedia of Food Microbiology: Second Edition*, pp. 69–76. <https://doi.org/10.1016/B978-0-12-384730-0.00009-4>
- Gómez-Godínez, L. J., Aguirre-Noyola, J. L., Martínez-Romero, E., Arteaga-Garibay, R. I., Ireta-Moreno, J. and Ruvalcaba-Gómez, J. M., 2023. A Look at Plant-Growth-Promoting Bacteria. *Plants*, 12(8), p. 1668. <https://doi.org/10.1590/S1415-475738420150053>
- Guo, L., Zheng, S., Cao, C. and Li, C., 2016. Tillage practices and straw-returning methods affect topsoil bacterial community and organic C under a rice-wheat cropping system in central China. *Scientific Reports*, 6(1), 33155. <https://doi.org/10.1038/srep33155>
- Hatten, J. and Liles, G., 2019. A ‘healthy’ balance – The role of physical and chemical properties in maintaining forest soil function in a changing world. *Developments in Soil Science*, 36, pp. 373–396. <https://doi.org/10.1016/B978-0-444-63998-1.00015-X>
- Hernández, M., Dumont, M. G., Yuan, Q., and Conrad, R., 2015. Different bacterial populations associated with the roots and rhizosphere of rice incorporate plant-derived carbon. *Applied and Environmental Microbiology*, 81(6), pp. 2244–2253. [https://doi.org/10.1128/aem.03209-14/suppl\\_file/zam999116128so1.pdf](https://doi.org/10.1128/aem.03209-14/suppl_file/zam999116128so1.pdf)
- Hu, J., Zhao, Y., Yao, X., Wang, J., Zheng, P., Xi, C. and Hu, B., 2021. Dominance of comammox *Nitrospira* in soil nitrification. *Science of The Total Environment*, 780, p. 146558. <https://doi.org/10.1016/J.SCITOTENV.2021.146558>
- Junior, D.B., Guarçoni, R.C., da Silva, M.D.C.S., Veloso, T.G.R., Kasuya, M.C.M., da Silva Oliveira, E.C., da Luz, J.M.R., Moreira, T.R., Debona, D.G. and Pereira, L.L., 2021. Microbial fermentation affects sensorial, chemical, and microbial profile of coffee under carbonic maceration. *Food Chemistry*, 342, p. 128296. <https://doi.org/10.1016/j.foodchem.2020.128296>
- Kalra, Y.P., 1995. Determination of pH of soils by different methods: collaborative study. *Journal of AOAC International*, 78(2), p. 310–324. <https://doi.org/10.1093/jaoac/78.2.310>

- Kämpfer, P., Dreyer, U., Neef, A., Dott, W. and Busse, H. J., 2003. *Chryseobacterium defluvii* sp. nov., isolated from wastewater. *International Journal of Systematic and Evolutionary Microbiology*, 53, pp. 93–97. <https://doi.org/10.1099/IJS.0.02073-0>
- Kämpfer, P., Vaneechoutte, M., Lodders, N., De Baere, T., Avesani, V., Janssens, M., Busse, H.J. and Wauters, G., 2009. Description of *Chryseobacterium anthropi* sp. nov. to accommodate clinical isolates biochemically similar to *Kaistella koreensis* and *Chryseobacterium haifense*, proposal to reclassify *Kaistella koreensis* as *Chryseobacterium koreense* comb. nov. and emended description of the genus *Chryseobacterium*. *International Journal of Systematic and Evolutionary Microbiology*, 59(10), pp. 2421–2428. <https://doi.org/10.1099/IJS.0.008250-0>
- Katsenios, N., Andreou, V., Sparangis, P., Djordjevic, N., Giannoglou, M., Chanioti, S., Kasimatis, C.N., Kakabouki, I., Leonidakis, D., Danalatos, N., Katsaseros, G. and Efthimiadou, A., 2022. Assessment of plant growth promoting bacteria strains on growth, yield and quality of sweet corn. *Scientific Reports*, 12(1), p. 11598. <https://doi.org/10.1038/s41598-022-16044-2>
- Kejela, T., Thakkar, V. R. and Thakor, P., 2016. *Bacillus* species (BT42) isolated from *Coffea arabica* L. rhizosphere antagonizes *Colletotrichum gloeosporioides* and *Fusarium oxysporum* and also exhibits multiple plant growth promoting activity. *BMC Microbiology*, 16(1), pp. 1–13. <https://doi.org/10.1186/S12866-016-0897-Y>
- Keren, R. 1996. Boron. In: D.L. Sparks, A.L. Page, R.H. Helmke, R.H. Loeppert, P.N. Soltanpour, M.A. Tabatabal, C.T. Johnston and M.E. Sumner, eds. *Methods of Soil Analysis: Part 3 Chemical Methods*, Madison: Soil Science Society of America, American Society of Agronomy, pp. 603–626. <https://doi.org/10.2136/sssabookser5.3.c21>
- Kuffner, M., Puschenreiter, M., Wieshammer, G., Gorfer, M. and Sessitsch, A., 2008. Rhizosphere bacteria affect growth and metal uptake of heavy metal accumulating willows. *Plant and Soil*, 304(1–2), pp. 35–44. <https://doi.org/10.1007/S11104-007-9517-9>
- Lahlali, R., Ezrari, S., Radouane, N., Kenfaoui, J., Esmaeel, Q., El Hamss, H., Belabess, Z. and Barka, E.A., 2022. Biological Control of Plant Pathogens: A Global Perspective. *Microorganisms*, 10(3), p. 596. <https://doi.org/10.3390/microorganisms10030596>
- Lian, T., Jin, J., Wang, G., Tang, C., Yu, Z., Li, Y., Liu, J., Zhang, S. and Liu, X., 2017. The fate of soybean residue-carbon links to changes of bacterial community composition in Mollisols differing in soil organic carbon. *Soil Biology and Biochemistry*, 109, pp. 50–58. <https://doi.org/10.1016/j.soilbio.2017.01.026>
- Lin, C. and Coleman, N.T. 1960. The measurement of exchangeable aluminum in soils and clays. *Soil Science Society of America Journal*, 24(6), pp. 444–446. <https://doi.org/10.2136/sssaj1960.03615995002400060009x>
- Lladó, S., López-Mondéjar, R. and Baldrian, P., 2017. Forest Soil Bacteria: Diversity, Involvement in Ecosystem Processes, and Response to Global Change. *Microbiology and Molecular Biology Reviews*, 81, p. e00063–16. <https://doi.org/10.1128/MMBR.00063-16>
- Lopes, M.J. dos S., Dias-Filho, M.B. and Gurgel, E.S.C., 2021. Successful Plant Growth-Promoting Microbes: Inoculation Methods and Abiotic Factors. *Frontiers in Sustainable Food Systems*, 5, p. 48. <https://doi.org/10.3389/FSUFS.2021.606454/XML/NLM>
- López Báez, W., Castro Mendoza, I., Salinas Cruz, E., Reynoso Santos, R. and López Martínez, J., 2016. Properties of soils coffee in the Biosphere Reserve El Triunfo, Chiapas, Mexico. *Revista Mexicana de Ciencias Agrícolas*, 7(3), pp. 607–618.
- Mathew, R.P., Feng, Y., Githinji, L., Ankumah, R. and Balkcom, K.S., 2012. Impact of no-tillage and conventional tillage systems on soil microbial communities. *Applied and Environmental Soil Science*, 2012, p. 548620. <https://doi.org/10.1155/2012/548620>
- Maurya, A., Kesharwani, L. and Mishra, M.K., 2018. Analysis of heavy metal in soil through atomic absorption spectroscopy for forensic consideration. *International Journal for Research in Applied Science and Engineering Technology*, 6(6), pp. 1188–1192.

- Miller, J.A., Kalyuzhnaya, M.G., Noyes, E., Lara, J.C., Lidstrom, M.E. and Chistoserdova, L., 2005. *Labrys methylaminiphilus* sp. nov., a novel facultatively methylotrophic bacterium from a freshwater lake sediment. *International Journal of Systematic and Evolutionary Microbiology*, 55(3), pp. 1247–1253. <https://doi.org/10.1099/ijs.0.63409-0>
- Mussatto, S.I., Machado, E.M.S., Martins, S. and Teixeira, J.A., 2011. Production, Composition, and Application of Coffee and Its Industrial Residues. *Food and Bioprocess Technology*, 4(5), pp. 661–672. <https://doi.org/10.1007/S11947-011-0565-Z>
- Nakatsu, C.H., Hristova, K., Hanada, S., Meng, X.Y., Hanson, J.R., Scow, K.M. and Kamagata, Y., 2006. *Methylibium petroleiphilum* gen. nov., sp. nov., a novel methyl tert-butyl ether-degrading methylotroph of the Betaproteobacteria. *International Journal of Systematic and Evolutionary Microbiology*, 56(5), pp. 983–989. <https://doi.org/10.1099/ijs.0.63524-0>
- Poria, V., Dębiec-Andrzejewska, K., Fiodor, A., Lyzohub, M., Ajijah, N., Singh, S. and Pranaw, K., 2022. Plant Growth-Promoting Bacteria (PGPB) integrated phytotechnology: A sustainable approach for remediation of marginal lands. *Frontiers in Plant Science*, 13, p. 999866. <https://doi.org/10.3389/fpls.2022.999866>
- Pozzi, R., Simone, M., Mazzetti, C., Maffioli, S., Monciardini, P., Cavaletti, L., Bamonte, R., Sosio, M., and Donadio, S., 2011. The genus *Actinoallomurus* and some of its metabolites. *The Journal of Antibiotics*, 64(1), pp. 133–139. <https://doi.org/10.1038/ja.2010.149>
- Pukall, R., Lapidus, A., Glavina Del Rio, T., Copeland, A., Tice, H., Cheng, J. F., Lucas, S., Chen, F., Nolan, M., Bruce, D., Goodwin, L., Pitluck, S., Mavromatis, K., Ivanova, N., Ovchinnikova, G., Pati, A., Chen, A., Palaniappan, K., Land, M. and Hugenholtz, P., 2010. Complete genome sequence of *Conexibacter woesei* type strain (ID131577T). *Standards in Genomic Sciences*, 2(2), p. 212. <https://doi.org/10.4056/SIGS.751339>
- Rawat, S. R., Männistö, M. K., Bromberg, Y. and Häggblom, M. M., 2012. Comparative genomic and physiological analysis provides insights into the role of Acidobacteria in organic carbon utilization in Arctic tundra soils. *FEMS Microbiology Ecology*, 82(2), 341–355. <http://doi.org/10.1111/j.1574-6941.2012.01381.x>
- Rincon-Florez, V.A., Carvalhais, L.C., and Schenk, P.M., 2013. Culture-Independent Molecular Tools for Soil and Rhizosphere Microbiology. *Diversity*, 5(3), pp. 581–612. <https://doi.org/10.3390/D5030581>
- Salazar, S., Ochoa, A. and McCarthy, P.J., 2023. How the visit of pedologist Hans Jenny to Colombia (1946–1947) contributed to the theory of soil-forming factors. *Geoderma*, 437, p. 116575.
- Sanford, R.A., Cole, J.R. and Tiedje, J.M., 2002. Characterization and Description of *Anaeromyxobacter dehalogenans* gen. nov., sp. nov., an Aryl-Halo-respiring Facultative Anaerobic Myxobacterium. *Applied and Environmental Microbiology*, 68(2), 893. <https://doi.org/10.1128/AEM.68.2.893-900.2002>
- Saxena, A.K., Kumar, M., Chakdar, H., Anuroopa, N., and Bagyaraj, D.J., 2020. *Bacillus* species in soil as a natural resource for plant health and nutrition. *Journal of Applied Microbiology*, 128(6), pp. 1583–1594. <https://doi.org/10.1111/JAM.14506>
- Schaefer, D. 1973. *Beitraege zur Klassifizierung und Taxonomie der Actinoplanaceen*. Dissertation, Marburg.
- Schmidt, R., Gravuer, K., Bossange, A.V., Mitchell, J. and Scow, K., 2018. Long-term use of cover crops and no-till shift soil microbial community life strategies in agricultural soil. *PloS one*, 13(2), p. e0192953. <https://doi.org/10.1371/journal.pone.0192953>
- Seki, T., Matsumoto, A., Shimada, R., Inahashi, Y., Omura, S. and Takahashi, Y., 2012. *Conexibacter arvalis* sp. nov., isolated from a cultivated field soil sample. *International Journal of Systematic and Evolutionary Microbiology*, 62(10), pp. 2400–2404. <https://doi.org/10.1099/ijs.0.036095-0>
- Séneca, J., Pjevac, P., Canarini, A., Herbold, C.W., Zioutis, C., Dietrich, M., Simon, E., Prommer, J., Bahn, M., Pötsch, E. M., Wagner, M., Wanek, W. and Richter, A., 2020. Composition and activity of nitrifier communities in soil are unresponsive to

- elevated temperature and CO<sub>2</sub>, but strongly affected by drought. *The ISME Journal*, 14(12), pp. 3038–3053. <https://doi.org/10.1038/s41396-020-00735-7>
- Shen, F.T., Kämpfer, P., Young, C.C., Lai, W.A. and Arun, A.B., 2005. *Chryseobacterium taichungense* sp. nov., isolated from contaminated soil. *International Journal of Systematic and Evolutionary Microbiology*, 55(3), pp. 1301–1304. <https://doi.org/10.1099/ijs.0.63514-0>
- Souza, R.D., Ambrosini, A. and Passaglia, L.M., 2015. Plant growth-promoting bacteria as inoculants in agricultural soils. *Genetics and Molecular Biology*, 38, pp. 401–419. <https://doi.org/10.1590/S1415-475738420150053>
- Soylak, M., Karatepe, A.U., Elçi, L. and Doğan, M., 2003. Column preconcentration/separation and atomic absorption spectrometric determinations of some heavy metals in table salt samples using amberlite XAD-1180. *Turkish Journal of Chemistry*, 27(2), pp. 235–242. <https://journals.tubitak.gov.tr/chem/vol27/iss2/11/>
- Spain, A. M., Krumholz, L. R., and Elshahed, M. S., 2009. Abundance, composition, diversity and novelty of soil Proteobacteria. *The ISME Journal*, 3(8), pp. 992–1000. <https://doi.org/10.1038/ismej.2009.43>
- Sumner, M.E. and Miller, W.P. 1996. Cation exchange capacity and exchange coefficients. In: D.L. Sparks, A.L. Page, R.H. Helmke, R.H. Loeppert, P.N. Soltanpour, M.A. Tabatabal, C.T. Johnston and M.E. Sumner, eds. *Methods of Soil Analysis: Part 3 Chemical Methods*, Madison: Soil Science Society of America, American Society of Agronomy, pp. 1201–1229. <https://doi.org/10.2136/sssabookser5.3.c40>
- Sun, R., Li, W., Dong, W., Tian, Y., Hu, C. and Liu, B., 2018. Tillage changes vertical distribution of soil bacterial and fungal communities. *Frontiers in Microbiology*, 9, p. 699. <https://doi.org/10.3389/fmicb.2018.00699>
- Urgiles-Gómez, N., Avila-Salem, M. E., Loján, P., Encalada, M., Hurtado, L., Araujo, S., Collahuazo, Y., Guachanamá, J., Poma, N., Granda, K., Robles, A., Senés, C. and Cornejo, P., 2021. Plant growth-promoting microorganisms in coffee production: from isolation to field application. *Agronomy*, 11(8), p.1531. <https://doi.org/10.3390/agronomy11081531>
- Valencia-Cantero, E., Hernández-Calderón, E., Velázquez-Becerra, C., López-Meza, J.E., Alfaro-Cuevas, R. and López-Bucio, J., 2007. Role of dissimilatory fermentative iron-reducing bacteria in Fe uptake by common bean (*Phaseolus vulgaris* L.) plants grown in alkaline soil. *Plant and Soil*, 291(1–2), pp. 263–273. <https://doi.org/10.1007/s11104-007-9191-y/figures/5>
- Valinsky, L., Vedova, G. Della, Scupham, A.J., Alvey, S., Figueroa, A., Yin, B., Hartin, R.J., Chrobak, M., Crowley, D. E., Jiang, T. and Borneman, J., 2002. Analysis of bacterial community composition by oligonucleotide fingerprinting of rRNA genes. *Applied and Environmental Microbiology*, 68(7), pp. 3243–3250. <https://doi.org/10.1128/aem.68.7.3243-3250.2002>
- Velázquez-Becerra, C., Macías-Rodríguez, L.I., López-Bucio, J., Altamirano-Hernández, J., Flores-Cortez, I. and Valencia-Cantero, E., 2011. A volatile organic compound analysis from *Arthrobacter agilis* identifies dimethylhexadecylamine, an amino-containing lipid modulating bacterial growth and *Medicago sativa* morphogenesis in vitro. *Plant and Soil*, 339(1), pp. 329–340. <https://doi.org/10.1007/S11104-010-0583-Z>
- Veloso, T.G.R., da Silva, M.D.C.S., Cardoso, W.S., Guarçoni, R.C., Kasuya, M.C.M. and Pereira, L.L., 2020. Effects of environmental factors on microbiota of fruits and soil of *Coffea arabica* in Brazil. *Scientific Reports*, 10(1), p. 14692. <https://doi.org/10.1038/s41598-020-71309-y>
- Wang, K., Jia, R., Li, L., Jiang, R., and Qu, D., 2020. Community structure of Anaeromyxobacter in Fe(III) reducing enriched cultures of paddy soils. *Journal of Soils and Sediments*, 20(3), pp. 1621–1631. <https://doi.org/10.1007/s11368-019-02529-7/tables/2>
- Wang, Y., Li, C., Tu, C., Hoyt, G.D., DeForest, J.L. and Hu, S., 2017. Long-term no-tillage and organic input management enhanced the diversity and stability of soil microbial community. *Science of the Total Environment*, 609, p. 341–347. <https://doi.org/10.1016/j.scitotenv.2017.07.0>

## 53

- Williams, S.D., Barkla, B. J., Rose, T.J. and Liu, L., 2022. Does coffee have terroir and how should it be assessed? *Foods*, 11(13), p. 1907. <https://doi.org/10.3390/foods11131907>
- Wuenschel, R., Unterfrauner, H., Peticzka, R. and Zehetner, F., 2015. A comparison of 14 soil phosphorus extraction methods applied to 50 agricultural soils from Central Europe. *Plant, Soil and Environment*, 61(2), pp. 86-96. <http://doi.org/10.17221/932/2014-PSE>
- Xin, X.L., Yang, W.L., Zhu, Q.G., Zhang, X.F., Zhu, A.N. and Zhang, J.B., 2018. Abundance and depth stratification of soil arthropods as influenced by tillage regimes in a sandy loam soil. *Soil Use and Management*, 34(2), 286-296. <https://doi.org/10.1111/sum.12412>
- Yin, C., Casa Vargas, J.M., Schlatter, D.C., Hagerty, C.H., Hulbert, S.H. and Paulitz, T.C., 2021. Rhizosphere community selection reveals bacteria associated with reduced root disease. *Microbiome*, 9(1), pp. 1-18. <https://doi.org/10.1186/S40168-020-00997-5/tables/4>
- Yin, C., Jones, K.L., Peterson, D.E., Garrett, K.A., Hulbert, S.H. and Paulitz, T.C., 2010. Members of soil bacterial communities sensitive to tillage and crop rotation. *Soil Biology and Biochemistry*, 42(12), pp. 2111-2118. <https://doi.org/10.1016/j.soilbio.2010.08.006>
- Youssef, N.H., and Elshahed, M.S., 2008. Diversity rankings among bacterial lineages in soil. *The ISME Journal*, 3(3), pp. 305-313. <https://doi.org/10.1038/ismej.2008.106>
- Yu, Y., Lee, C., Kim, J. and Hwang, S., 2005. Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnology and Bioengineering*, 89(6), pp. 670-679. <https://doi.org/10.1002/bit.20347>
- Yunan, D., Xianliang, Q. and Xiaochen, W., 2018. Study on cation exchange capacity of agricultural soils. *IOP Conference Series: Materials Science and Engineering*, 392, p. 042039. <http://doi.org/10.1088/1757-899X/392/4/042039>
- Zhang, L., Gao, J.S., Kim, S.G., Zhang, C.W., Jiang, J.Q., Ma, X.T., Zhang, J. and Zhang, X.X., 2016. *Novosphingobium oryzae* sp. nov., a potential plant-promoting endophytic bacterium isolated from rice roots. *International Journal of Systematic and Evolutionary Microbiology*, 66(1), pp. 302-307. <https://doi.org/10.1099/ijsem.0.000718>
- Zheng, Y., Jin, J., Wang, X., Kopittke, P.M., O'Sullivan, J.B. and Tang, C., 2023. Disentangling the effect of nitrogen supply on the priming of soil organic matter: A critical review. *Critical Reviews in Environmental Science and Technology*, 54, pp. 676-697. <https://doi.org/10.1080/10643389.2023.2266312>
- Zhou, L.Y., Chen, X.Y., Du, Z.J., and Mu, D.S., 2019. *Pedobacter chinensis* sp. nov., a cellulose-decomposing bacterium from arctic tundra soil. *International Journal of Systematic and Evolutionary Microbiology*, 69(7), pp. 1926-1933. <https://doi.org/10.1099/ijsem.0.003403>