



## MOLECULAR CHARACTERIZATION AND IDENTIFICATION OF BIOCONTROL ISOLATES OF *Trichoderma harzianum* FROM EMBU DISTRICT, KENYA

### [CARACTERIZACIÓN MOLECULAR E IDENTIFICACIÓN DE AISLAMIENTOS DE BIOCONTROL DE *Trichoderma harzianum* DEL DISTRITO DE EMBU, KENIA]

E. N. Siameto<sup>1</sup>, S. Okoth<sup>2\*</sup>, N. O. Amugune<sup>2</sup>, and N. C. Chege<sup>2</sup>

<sup>1</sup> School of Sciences, Narok University College, P.O Box 861, Narok

<sup>2</sup> School of Biological Sciences, University of Nairobi, P.O Box 30197, Nairobi  
E-mail: dorisokoth@yahoo.com Tel: 02-444904 Ext 2483.

\*Corresponding author

#### SUMMARY

Species in the genus *Trichoderma* are important commercial source of several enzymes, biofungicides and growth promoters. The most common biological control agents of the genus are strains of *T. harzianum*, *T. viride* and *T. viriens*. In this study, sixteen selected isolates of *T. harzianum* from different land use types in Embu, Kenya were tested for antagonistic action against five soil borne phytopathogenic fungi (*Rhizoctonia solani*, *Pythium sp.*, *Fusarium graminearum*, *F. oxysporum f. sp. phaseoli* and *F. oxysporum f. sp. Lycopersici*) using dual culture assay and through production of non-volatile inhibitors. Seven isolates were further characterized using RAPD-PCR procedure to determine genetic variability. All *T. harzianum* isolates had considerable antagonistic effect on mycelial growth of the pathogens in dual cultures compared to the control. Maximum inhibitions occurred in *Pythium sp.*-055E interactions (73%). The culture filtrates obtained from Czapek's liquid medium reduced the dry weight (mg) of the mycelia significantly while those from the potato dextrose broth showed minimum inhibition growth. *Pythium sp.* was most sensitive compared to other pathogens. Genetic similarities generated using Jaccard's coefficient of similarity ranged from 0.231 to 0.857 for isolates 055E, 011E, 010E and 015E. Since all *T. harzianum* isolates evaluated were effective in controlling colony growth of the soil borne pathogens both in dual cultures and in culture filtrates they could be tried as a broad spectrum biological control agent in the green house and under field conditions.

**Key words:** *Trichoderma harzianum*, growth antagonism, genetic similarity RAPDs

#### INTRODUCTION

*Trichoderma harzianum* is an asexually reproducing filamentous fungus and a species aggregate. It is grouped on the basis of conidiophores branching

patterns with short side branches, short inflated phialides, smooth and small conidia (Rifai 1969). These characteristics allow for the relatively easy identification of *Trichoderma* as a genus, but the species concept is difficult to interpret and there is considerable confusion over the application of specific names (Rifai 1969; Papavizas 1985). This disparity of criteria makes it difficult to search for, and above all characterize new biocontrol agents (BCAs) within the species and to reidentify them in natural environment once they are present in a pathosystem (Papavizas 1985).

The advent of polymerase chain reaction (PCR) has allowed the analysis of small numbers of fungal cells or even single spores, dried herbarium material (Carlies & Watkinson 1994), or extinct organisms (Golenberg *et al.* 1990). The selection of universal oligonucleotide primers specific to fungi (Van Belkum *et al.* 1993; Sandhu *et al.* 1995) has provided easy access to nucleotide sequences. These techniques have been proven to be valuable tools in fungal taxonomy and their application has led to the reconsideration of several genera (Sherriff *et al.*, 1994)

Hence, Random markers as products of the PCR-RAPD technique have been used for taxonomy of *Trichoderma* to discriminate species (Hadrys *et al.*, 1992; Williams *et al.*, 1990).

*Trichoderma* species have been investigated for over 70 years (Hjeljord & Tronsmo 1998). They have been used as biological control agents (BCAs) and their isolates have become commercially available of late (Freeman *et al.*, 2004). This development is largely the result of a change in public attitude towards the use of chemical pesticides and fungicides such as methyl bromide (Elad *et al.* 1980; Basim *et al.* 1999). In this respect *Trichoderma* species have been studied as BCAs against soil-borne plant pathogenic fungi (Henis 1984; Chet and Inbar, 1994). Replacement or reduction of chemical application can be achieved through use of biologically based fungicides, a concept

included in the broad definition of biological control proposed by Cook & Baker (1983). The commercial use of *Trichoderma* BCAs must be preceded by precise identification, adequate formulation, and studies about the synergistic effects of their mechanisms of biocontrol.

In this study the *in vitro* antifungal activity of *T. harzianum* isolates were evaluated against selected soil borne plant pathogens (*Rhizoctonia solani*, *Pythium sp.*, *Fusarium graminearum*, *F. oxysporum f. sp. phaseoli* and *F. oxysporum f. sp. Lycopersici*). We aimed also at characterizing *T. harzianum* isolates using RAPD-PCR technique to establish the degree of genetic variation and determine any relationship between molecular variation and antifungal activity.

## MATERIAL AND METHODS

### Isolation and identification of *Trichoderma* species

*Trichoderma sp.* were isolated using dilution plate technique (Johnson *et al.*, 1959) and soil washing methods (Gams *et al.*, 1987; Bills & Polishook 1994) on malt extract agar (MEA) and cornmeal agar (CMA) with 2% dextrose) both with streptomycin 50mg/L and cyclosporin 10mg/L. The colonies were counted and identified and transferred to Petri dishes containing potato dextrose agar (PDA) and incubated at 15, 25, 30 and 35°C for further identification to species level.

Genus identification of green fungus was undertaken using the method of Domsch *et al.* (1980). *Trichoderma* isolates were identified to species level following the taxonomic key of Samuels *et al.* (2004). Microscopic examination was carried out by mounting the culture in lactophenol cotton blue but for size measurements KOH and water was used as the mounting fluid. A small amount of material was placed in a drop of 3% KOH on a slide and then replaced with water. The isolates were preserved in sterile soil and stored at 4°C.

### Isolation of phytopathogenic fungi.

Diseased plant tissues were obtained from the University of Nairobi, College of Agriculture and Veterinary Medicine (CAVM), Upper Kabete campus field station which is 10 Km north-west of Nairobi. *Rhizoctonia solani* was isolated from Spinach (*Spinacea oleracea*), *Fusarium oxysporum f. sp. Lycopersici* from tomato (*Lycopersicum esculentum*), *F. oxysporum f. sp. phaseoli* from beans (*Phaseolus vulgaris*) and *F. graminearum* from maize cob (*Zea mays*). Diseased plant tissues were washed under running tap water to remove surface soil, dust and other contaminants. Tissue pieces were cut out from the leading edge of lesion, and placed in one percent

sodium hypochlorite for five minutes, then washed in sterile distilled water and dried on sterile filter paper. The dried pieces were cut into approximately one centimeter pieces, plated onto PDA and incubated at 25°C. *Pythium sp.* was obtained by planting beet root seeds in water logged soil. The seedlings that emerged were infected by dumping- off. The seedlings stems were cut into one centimeter pieces and inoculated onto PDA plates.

### Evaluation of Dual culture on agar plates

Plates of PDA were inoculated with five millimeter disc obtained from five-day-old cultures of the phytopathogens ten millimeter from the edge of the plate. After two days a five millimeter disc of the *Trichoderma harzianum* cultures was placed in the same plate at a distance of fifty five millimeter from the phytopathogens disc. *Pythium sp.* and the biocontrol strains were inoculated at the same time. The paired cultures were incubated at room temperature for six days. The growth of the fungi was recorded by measuring the radial growth of the pathogens. The percentage growth of the pathogens was calculated as follows:

Percent growth =  $\frac{\text{Radius of the growth in the direction of the test strain}}{\text{radius of the growth in the absence of the test strain}} \times 100$  (Edington *et al.*, 1971). -----  
Equation 1

### Evaluation of dual culture using Slide method

For *Rhizoctonia solani* and *Pythium sp.*-*Trichoderma* interaction, a clean sterile glass slide was placed in nine centimeter diameter plates. A small amount of melted PDA was spread over the slide to make a thin film on the slide. Five millimeter discs of one week old growing colonies cut from the margin of each pathogen and *Trichoderma* isolates were placed on the opposite sides of the slide three centimeter apart on the PDA surface. A two milliliter of sterilized distilled water was added to the plate to prevent drying and incubated  $25 \pm 1^\circ\text{C}$  for a week. At the end of incubation period, point of contact between *Trichoderma*-Pathogen hyphae was stained with lactophenol in cotton blue and observed under a light microscope for the presence of mycelial penetration and for cell wall disintegration.

### Determination of antifungal properties of *T. harzianum* culture filtrates

To harvest non-volatile antibiotics produced by *Trichoderma* isolates, the fungus was grown on potato dextrose broth (PDB) and Czapek's liquid medium (CLM)

Each *T. harzianum* isolate was inoculated into 100ml PDB incubated at 20°C then filtered through 0.22 mm Millipore filters after 10 days. 2ml aliquots of these filtrates were placed in sterile Petri dishes and 25ml of PDA added. Five millimeter wide mycelial discs of the pathogen were placed at the center of the solidified agar plates and incubated at room temperature for 6 days. Colony diameters were measured daily and inhibition percentage obtained using the formula:

Inhibition percent =  $[(C2-C1) \div C2] \times 100$  (Edington *et al.* 1971). -----Equation 2

Where: C1 means growth of the pathogens in the presence of antagonist and C2 means the growth of control. Each experiment had three replicates and complete randomized design was adopted. PDA without the culture filtrates served as the control.

To test for non-volatile antibiotics from Czapek's liquid medium the method of Tianhui (1994) was amended. Flasks containing 50 ml of CLM were inoculated with a five millimeter disc of *T. harzianum* isolates and incubated at 25°C in the dark for two weeks. Culture liquids were filtered through filter paper and sterilized by bacterial filtration (Acrodisc nitrocellulose membranes, pore size 0.22µm). The filtrates were diluted with PDB to a concentration of 70%. PDB lacking the filtrate of *Trichoderma* served as a control. Two agar discs containing the pathogen were inoculated in each flask and then incubated at 25°C and mycelia filtered through filter paper after eight days, dried and weighed. The percent dry weight was then obtained using the following formula:

Percent dry weight =  $\frac{\text{dry weight of control} - \text{dry weight of pathogen}}{\text{dry weight of control}} \times 100$ . -----Equation 3.

Each treatment was replicated three times.

### Isolation of DNA from filamentous fungi

Mycelia for DNA extraction were cultured in 50 ml of yeast potato dextrose broth (YPDB) at 20°C with rotary shaking at 120 revolutions per minute (rpm), harvested by filtration through a filter paper and washed with distilled water after two days. The mycelia were freeze-dried and ground in the presence of sand and stored at -20°C. One hundred milligrams of the mycelial powder was transferred into eppendorf tubes. 500µl of 2xCTAB buffer (equilibrated to 65°C) and 1.0% of β-mercaptoethanol were added and the tubes heated at 65°C for 30 minutes. 500µl of chloroform-isoamyl alcohol (24:1) was added to the tubes and then vortexed for 30 seconds then centrifuged at 12,000rpm for 15 minutes. The upper portion of the aqueous phase was recovered and 1/5

volume of 5% CTAB and mix. Another chloroform-isoamyl alcohol (24:1) extraction was performed. Equal volume of CTAB precipitation buffer was mixed with the recovered supernatant and left to stand on ice for 20 minutes. The mixture was centrifuged for 15 minutes at 12,000rpm and the supernatant discarded. The pellet was rehydrated with high salt TE buffer (heat at 65°C for 5-10 minutes). DNA was precipitated by the addition of an equal volume of cold absolute ethanol with incubation at -20°C for 10 min. DNA was collected by centrifugation at 12,000rpm for 10 minutes, washed with 70% ethanol, dried, and re-suspended in 100µl of 1XTE buffer and stored at -20°C. The quality of the DNA was checked by use of 0.7% agarose gel and quantified spectrophotometrically.

### RAPD analysis and PCR conditions

Amplification reactions were performed in 0.6ml microcentrifuge tubes in a 25 µl reaction volumes containing 5ng of template DNA, *Taq* buffer, 2.5mM MgCl<sub>2</sub>, 0.8mM each dNTP, 1.15ng/µl Primer and 1unit *Taq* DNA polymerase. Amplification reactions were performed in a Perkin-Elmer, Gene amp PCR system 2400 thermal Cycler programmed for 35 cycles of denaturation at 94°C for 30 seconds, low stringency annealing temperature at 31°C for 1.0 minute and polymerization at 72°C for 1.0 minute with a final extension step at 72°C for 10 minutes. PCR products were separated on 2.5 % agarose/ 1X TBE gels. A 50bp DNA molecular size marker was loaded on the first well and used for comparison. The banding pattern was visualized on Ultraviolet transilluminator and documented by MultiDoc digital imaging system.

### Data analysis

Bands were manually scored 1 for presence and 0 for absence and the binary data used for statistical analysis using the software R-command version 2.1.1. The size of the fragments (molecular weight in base pairs) was estimated by using 50 bp ladder marker, which was run along with the amplified products. A genetic dissimilarity matrix was calculated according to Squared Euclidean Distance which estimated all pairwise differences in the amplification product and Cluster analysis was done by Wards method using a minimum variance algorithm (Ward, 1963).

Genetic Similarity (GS) was analyzed using the equation (Jaccard, 1908):

$GS = (N_{ab}) / (N_a + N_b - N_{ab})$  ----- Equation 4.

Where  $N_{ab}$  is the number of shared fragments between isolates a and b,  $N_a$  is the number of scored fragments of isolate a, and  $N_b$  the number of scored fragments of isolate b. Genetic distance (GD) was then calculated as

GD = 1-GS. The data for growth inhibition measurement were arcsine transformed then subjected to analysis using the Statistica version 6 software.

## RESULTS

### Inhibition of growth of pathogenic fungi by *T. harzianum* on culture media

As single cultures *Pythium* sp grew actively and colonized the entire agar surface within two days, whereas *F. oxysporum* took two weeks, and *F. graminearum* and *Rhizoctonia* sp filled the plate in six days. Dual culture assays provided evidence that *T. harzianum* isolates reduced growth of the pathogens. Nine out of the sixteen isolates tested were able to inhibit the growth of three pathogens each by more than 50%. Three of the *T. harzianum* isolates inhibited four pathogens. Isolates 015E and 051E were superior to others since they inhibited the growth of five pathogens tested.

*T. harzianum* showed parasitic behavior against *Pythium* (Fig 1) by coiling round the host hyphae and degrading it. The dual cultures plates showed initial rapid growth of the host which stopped at the point of contact with the parasite. *T. harzianum* over grew the pathogen resulting into complete degradation of the latter and sporulation of the former over the entire plate. The isolate 055E gave the lowest percentage inhibition of 31.76% against *T. harzianum* while 015E gave the highest inhibition of 73.33 %.




Figure 1. Light microscope micrograph showing *Trichoderma harzianum* encircling mycelia of *Pythium* sp as shown by the arrows.

All isolates of *T. harzianum* that were tested inhibited growth of *R. solani* with isolate 029E giving the highest percentage inhibition (61.55 %) while 063E the lowest (25.88%). The inhibition process was parasitic (Fig. 2), where the parasite penetrated the host cell wall directly suggested that there was some mechanical activity that led to the disintegration of the hyphae (Fig 3).




Figure 2. Micrograph of light microscope showing coiling of *T. harzianum* over *R. solani* Arrows point to interaction zones between *T. harzianum* and *R. solani*.




Figure 3. Light microscope micrograph showing disintegrated hyphae of *R. solani*. The arrows indicate the disintegrated hyphae.

### Inhibition of growth of pathogenic fungi by *T. harzianum* through production of non-volatile compounds

Slight colony inhibition of the pathogens (*F. oxysporum* f.sp *Lycopersici*, *F. oxysporum* f. spp. *phaseoli*, *Pythium* spp., *Fusarium graminearum* and *Rhizoctonia solani*) was observed when exposed to the culture filtrates of *T. harzianum* (Table 1). There was no inhibition of radial growth by all tested isolates on *Pythium* sp. Isolate 014E inhibited *F. oxysporum* growth by 80%. Isolates 057E, 021E, 055E, 015E, 0051E, and 010E had no effect on *F. graminearum*. The culture filtrates from isolates 042E, 063E, 030E, 029E, 046E, 044E, 011E, 049E, and 014E inhibited the growth of *R. solani*.

The different culture filtrate from the *T. harzianum* isolates caused significant ( $P = 0.05$ ) effect on the dry weight of the phytopathogens: *F. oxysporum* f. sp *Lycopersici*, *F. oxysporum* f. sp *phaseoli* *Pythium* sp, *Fusarium graminearum* and *Rhizoctonia solani*. The culture filtrates of the isolates suppressed the growth of the pathogens and reduced the dry weight (mg)

mycelia (Fig 4). *Pythium sp* was the most influenced as compared to the control. However some isolates' filtrates seems to have enhanced the accumulation of mycelia since the percentage weight obtained of the mycelia was more than 100%. *Pythium sp* for example was enhanced by the isolates 042E, 049E, 046E, 045E, 021E and 051E while the rest (014E, 029E, 010E, 044E, 055E, 011E, 015E, 057E, and 031E) greatly suppressed the accumulation of the mycelia of *Pythium* by more than 50%. The filtrate from the isolate 011E was superior to others since the dry weight of the mycelia of four (*F. oxysporum f. sp Lycopersici*, *F. oxysporum f. sp phaseoli* *Pythium sp* and *F. graminearum*) phytopathogens was less than 50%. The *Fusarium* pathogens were always susceptible to the culture filtrates from all the *T. harzianum* isolates.

#### DNA polymorphism analysis of the *T. harzianum* isolates

In the study four random primers namely 203, 230, 220, 0p13 were used which gave bands ranging from 350bp to 2000bp as shown in Table 2. All the primers produced intense bands totaling to 81 bands were produced. The 7 samples used for DNA polymorphism were antagonistic to the phytopathogens used. In dual cultures the isolates 015E and 051E were superior to others since they inhibited the growth of five pathogens by more than 50% and 029E inhibited the growth of four pathogens. In the culture filtrate activity the isolates 011E and 055E had a wide range of activity by suppressing the growth of five pathogens while isolates 044E and 010E suppressed the growth of four pathogens each.

Table 1. Effect of different isolates of *T. harzianum* on radial growth inhibition (%) of phytopathogens by the production of non-volatile compounds from Potato Dextrose Broth after 6 days of incubation.

<i>T. harzianum</i> Isolates	Radial Growth Inhibition (%)				
	<i>F. oxysporum f.sp Lycopersici</i>	<i>F.oxysporum f. sp phaseoli</i>	<i>Pythium sp,</i>	<i>F. graminearum</i>	<i>R. solani</i>
042E	12.74	9.61	0	66.27	4.31
063E	2.27	13.46	0	44.70	17.64
030E	20.29	23.71	0	40.39	28.627
029E	32.51	1.92	0	85.88	11.76
046E	31.94	23.07	0	43.92	31.37
044E	40.08	20.51	0	36.47	30.98
057E	11.02	17.30	0	0	0
045E	19.72	27.56	0	6.66	0
011E	16.80	28.20	0	7.05	8.23
049E	4.02	10.89	0	5.88	5.88
021E	15.07	14.74	0	0	0
055E	14.50	26.28	0	0	0
015E	10.33	16.66	0	0	0
051E	11.55	15.38	0	0	0
010E	11.57	17.30	0	0	0
014E	80.22	82.05	0	8.62	71.37

Table 2. Synthetic deoxyribonucleotides used as primers for amplification of *Trichoderma harzianum* DNA.

Primer code	Primer sequence 5'- 3'	Total No. of bands	Range sizebands(bp)
203	CACGGCGAGT	7	600-2000
220	TCGATGTCG	7	600-2000
0P13	CAGCACCCAC	7	300-650
230	CGTCGCCCAT	7	350-1800




Figure 4. Percentage dry weight (mg) of mycelia of fungal phytopathogens p1: *F. oxysporum f. sp Lycopersici*, p2: *F. oxysporum f. sp phaseoli* p3: *Pythium sp.*, p4: *Fusarium graminearum* and p5: *Rhizoctonia solani* as influenced by non-volatile antibiotics of *T harzianum* isolates.

The DNA profiles obtained for *Trichoderma harzianum* isolates were scored (Fig 5) and a Dendrogram or Dissimilarity matrix developed using Squared Euclidean Distance and Clustering based on Wards method (Fig 6).




Figure 5. **A** RAPD profiles obtained with primer 203. Positions 2-9 corresponds to isolates T010, T015, T011, 021E, 051E, 029E, 055E, and 044E. **B** RAPD profiles obtained with primer 230. Positions 2-9 corresponds to isolates 044E, T011, T015, T014, 051E, 055E, T010, and 029E.




Figure 6. Dendrogram for 7 *Trichoderma harzianum* isolates as revealed by RAPD markers. Where : 1:044E, 2:011E, 3:015E, 4: 051E, 5: 055E, 6:010E, 7:029E

In the Dendrogram, all the 7 isolates were distinctly divided into two major clusters 'A' and 'B' at 20 units. Isolate 051E and 029E spanned the extremes of the entire Dendrogram. Genetic dissimilarity ranged from a lowest of 0.143 (between T010 and T015) to a highest of 0.857 (between 055E and 051E). Isolate 051E, T011, T015, and T010 were assigned to cluster 'A'. Genetic dissimilarity among the entries in this cluster ranged from a lowest of 14.3 percent (between T015 and T010) to a highest of 35.7 percent (between T010 and 051E). The other cluster 'B' comprised of three accessions. In this cluster isolate 044E 055E and 029E were grouped together. The genetic dissimilarity in this group was ranging from 33.3 percent between 055E and 029E to a high of 75 percent between 044E and 029E.

## DISCUSSION

*T. harzianum* inhibited the growth of the target organisms through its ability to grow much faster than the pathogenic fungi thus competing efficiently for space and nutrients. Starvation was the most common cause of death for microorganisms, so that competition for limiting nutrients resulted in biological control of fungal phytopathogens.

A second mechanism of pathogen control was mycoparasitism. Microscopic observation of the interaction region between *R. solani* and *Pythium* spp. with *T. harzianum* showed that the mycelia of *T. harzianum* grew on the surface of the pathogens always coiling round their mycelia and later penetrating their cell walls directly without formation of appressorium structures. The pathogen mycelia then disintegrated suggesting enzyme action. Lorito *et al.* (1998), Metcalf *et al.* (2001) and Sharon *et al.*, 2001 demonstrated possible role of chitinolytic and/or glucanases enzymes in bio-control by *Trichoderma*. These enzymes function by breaking down the polysaccharides, chitin, and glucans that are responsible for the rigidity of fungal cell walls, thereby destroying cell wall integrity limiting the growth of the pathogen. A mixture of several enzymes might be necessary for efficient cell wall lysis. *T. harzianum* has been reported to apply high chitinase and  $\beta$ -1, 3-glucanase activities (Sivan *et al.*, 1984).

A third mechanism of pathogen control by *Trichoderma* was through antibiosis. Culture filtrates of the isolates had an inhibitory effect on the radial growth of the pathogens and mycelial accumulation suggesting action of non-volatile antibiotics in the filtrates. This agrees with the findings of Sivan *et al.*, (1984) who noted that culture filtrates of *T. harzianum* strongly inhibited the growth of *Pythium aphanidermatum* whereas *T. hamatum* filtrates caused only minor inhibition of growth. Antibiotic inhibitions have been documented by Claydon *et al.* (1987),

Dubey and Suresh (2006), Kucuk and Kivanc (2003) and Lynch (1990). Claydon *et al.* (1987) reported inhibition due to antibiotics trichodermin, harzianum A and harzianolide. Dubey and Suresh (2006) found that non-volatile substances produced by *T. harzianum* are inhibitory to *F. oxysporum f. sp. ciceris* causing chickpea wilt.

Culture filtrate from PDB inhibited *Fusarium* spp. and *R. solani* but not *Pythium* sp. The difference in activity of the culture filtrates displayed by PDB and Czapek's liquid media on the pathogens indicates the importance of substrate in fungal production of secondary metabolites.

The presence of inhibition zones in dual cultures between *F. oxysporum f. sp. Lycopersici* and *T. harzianum* suggested secretion of diffusible non-volatile inhibitory substance by the *T. harzianum* isolates, which has also been documented by Grodona *et al.*, (1997) and Behzad *et al.*, (2008).

The DNA analysis of *T. harzianum* isolates from Embu showed existence of intraspecific variation. The genetic variation shown by the four random primers 203, 230, 220, 0p13 resulted in bands ranging from 350bp to 2000bp. This agrees with Moller *et al.* (1995) who detected intraspecific diversity not only between isolates of *Chaunopycnis alba* from different geographic regions or hosts, but also between isolates from a single location.

Based on RAPD data isolates from the same land use appeared in different groups due to intraspecific variation, for example isolate 044E and isolate 011E collected from camphor plantation were clustered in different groups and were 54.4% dissimilar. Goes *et al.*, (2002) also found intraspecific genetic variation among *Trichoderma* isolates. In his work the isolate Tm isolated from corn seed was grouped in the Dendrogram with the isolate 2820 isolated from sugar cane.

Table 3. Genetic distance among 7 *Trichoderma harzianum* isolates

	044E	011E	015E	051E	055E	010E	029E
044E	0						
011E	0.544	0					
015E	0.643	0.231	0				
051E	0.544	0.333	0.231	0			
055E	0.667	0.769	0.643	0.857	0		
010E	0.643	0.231	0.143	0.357	0.643	0	
029E	0.75	0.643	0.533	0.733	0.333	0.533	0

## CONCLUSION

*T. harzianum* is a good candidate for biological control due to the different modes of action the fungus employs in inhibiting the growth of other fungi. Through RAPD technique intraspecific genetic variation between *T. harzianum* isolates was observed. The results presented here support the existence of cryptic species with similar morphology.

## ACKNOWLEDGEMENTS

We would like to express our special thanks to the University of Nairobi for laboratory equipment and enabling environment to conduct research and the Conservation and sustainable management of below ground biodiversity (CSM-BGBD) project. A project executed by TSBF/CIAT with cofinancing from the Global Environment Facility (GEF) and implementation support from the United Nations

Environment Programme (UNEP) for financial support of the research.

## REFERENCES

- Basim, H., Ozturk, S.B. and Yegen, O. 1999. Efficacy of a biological fungicide (Planter Box *Trichoderma harzianum* Rifai T-22) against seedling root rot pathogens (*Rhizoctonia solani*, *Fusarium* sp) of cotton. GAP-Environmental Symposium. Sanliurfa. Turkey, 137-144.
- Behzad, H., Mousa, T., Mohammad, R.M. and Mahdi, D. 2008. Biological potential of some Iranian *Trichoderma* isolates in the control of soilborne plant pathogenic fungi. African Journal of Biotechnology. 7: 967-972.



- Bills, G. F. and Polishook, J. D. 1994. Abundance and diversity of microfungi in leaf litter of a lowland rain forest in Costa Rica. *Mycologia*. 86: 187 – 198.
- Carlie, M. J. and Watkinson, S. C. 1994. *The fungi*. Academic press, Ltd., London, United Kingdom.
- Chet, I. and Inbar, J. 1994. Biological control of fungal pathogens. *Applied Biochemistry and Biotechnology* 48: 37-43.
- Claydon, N., Allan, M., Hanso, J. R. and Avent, A. G. 1987. Antifungal alkyl pyrones of *Trichoderma harzianum*. *Transactions British Mycological Society*. 88: 503-513.
- Cook, R. J. and Baker, K. F. 1983. *The Nature and Practise of Biological Control of Plant Pathogens*. American Phytopathological Society, St Paul M N, 539.
- Domsch, K.H., Gams, W. and Anderson, T.H. 1980. *Compendium of soil fungi*. Academic Press London. 809.
- Dubey, S.C. and Suresh, M. 2006. Randomly Amplified Polymorphic DNA Markers for *Trichoderma* species and Antagonism against *Fusarium oxysporum f. sp. ciceris* Causing Chickpea Wilt. *Journal Phytopathology*. 154: 663-669.
- Edington, L.L., Khew, K.L. and Barron, G.I. 1971. Fungitoxic spectrum of benzimidazole compounds. *Phytopathology*. 61: 42-44.
- Elad, Y., Chet, I. and Katan, Y. 1980. *Trichoderma harzianum* a biocontrol agent effective against *Sclerotium rolfisii* and *Rhizoctonia solani*. *Phytopathology*. 70: 119-121.
- Freeman, S., Minz, D., Kolesnik, I., Barbul, O., Zreibil, A., Maymon, M., Nitzani, Y., Kirshner, B., Rav-David, D., Bilu, A., Dag, A., Shafir, S. and Elad, Y. 2004. *Trichoderma* biocontrol of *Colletotricum acutatum* and *Botrytis cinerea*, and survival in strawberry.
- Gams, W., Aa, H.A. van der; Plaats-Niterink, A.J. van der; Samson, R. A. and Stalpers, J. A. 1987. *CBS course of mycology*. Baarn, Netherlands; Centraalbureau voor Schimmelcultures. 3, 136
- Goes, L.B., Costa, A.B.L., Freire, L.L.C. and Oliveria, N.T. 2002. Randomly amplified polymorphic DNA of *Trichoderma* isolates and antagonism against *Rhizoctonia solani*. *Brazilian Archives Biology Technology*. 45: 151-160.
- Golenberg, E.M., Giannasi, D.E., Clegg, M.T., Smiley, C.J., Durbin, M., Henderson, D. and Zurawski, G. 1990. Chloroplast DNA sequence from a miocene *Magnolia* species. *Nature* 344: 656-658.
- Grondona, I., Hermosa, M.R., Tejada, M., Gomis, M. D., Mateos, P.F., Bridge, P., Monte, E. and García-Acha, I. 1997. Physiological and biochemical characterization of *Trichoderma harzianum*, a biological control agent against soilborne fungal plant pathogens. *Appl. Environ. Microbiol.* 63: 3189-3198.
- Hadrys, H., Balick, M. and Schierwater, B. 1992. Application of random amplified polymorphic DNA (RAPD) in molecular ecology. *Molecular Ecology*. 1: 55-63.
- Henis, C. 1984. Biological control. In: *Current perspectives in microbial ecology* Klug, M.J. Reddy, C. A. (Eds). American Society of Microbiology, Washington. 353-361.
- Hjeljord, L.G. and Tronsmo, A. 1998. *Trichoderma* and *Gliocladium* in biocontrol: an overview. In Kubicek, C.P. Harman, G.E. (Eds), *Trichoderma and Gliocladium*. Taylor & Francis, Ltd., London. United Kingdom. 135-151
- Jaccard, P. 1908. Nouvelles recherches sur la distribution florale. *Bul. Soc. Vaudoise Sci. Nat.* 44: 223-270.
- Johnson, L.E., Bond, C.J. and Fribourg, H. 1959. *Methods for studying soil microflora-plant disease relationships*. Minneapolis: Burgess Publishing Company.
- Kucuk, C. and Kivanc, M. 2003. Isolation of *Trichoderma* Spp. and determination of their antifungal, biochemical and physiological features. *Turkish Journal Biology*. 27: 247-253.
- Lorito, M., Woo, S.L., Garcia Fernandez, I., Colucci, G., Harman, G.E., Pintor-Toro, J.A., Filippone, E., Mucciflora, S., Lawrence, C.B., Zoina, A., Tuzun, S. and Scala, F. 1998. Genes from mycoparasitic fungi as a source for improving plant resistance to fungal pathogens. *Proceedings National Academy Science, USA*. 95: 7860-7865.

- Lynch, J.M. 1990. Fungi as antagonists. In: "New Directions in Biological Control: Alternative for suppressing Agricultural Pests and Diseases". Liss, New York. 243-253.
- Metcalf, D.D. and Wilson, C.C. 2001. The process of antagonism of *Sclerotium cepivorum* in white rot affected onion roots by *Trichoderma koningii*. *Plant Pathology*. 50: 249-257.
- Möller, C., Bühler, T. and Dreyfuss, M.M. 1995. Intraspecific genetic diversity of *Chaunopycnis alba* detected by random amplified polymorphic DNA assay. *Mycology Research* 99: 681-688.
- Papavizas, G.C. 1985. *Trichoderma* and *Gliocladium*: biology, ecology, and potential for biocontrol. *Annual Review Phytopathology*. 23: 23-54.
- Rifai, M.A. 1969. A revision of the genus *Trichoderma*. *Mycology*. 116: 1-56.
- Samuels, G. J., Chaverri, P., Farr, D.F. and McCray, E.B. 2004. USDA, Beltsville, USA. *Trichoderma* online systematic Botany and Mycology Laboratory, ARS, USDA. Retrieved September 20, 2004, from <http://nt.arsgrin.gov/taxadescriptions/keys/TrichodermaIndex.cfm>
- Sandhu, G.S., Kline, B.C., Stockman, L. and Roberts, G.D. 1995. Molecular probes for diagnosis of fungal infections. *Journal Clinical Microbiology*. 35: 2270-2274.
- Sherriff, C., Whelan, M.J., Arnold, G.M., Lafai, J.F., Brygoo, Y. and Bailey, J.A. 1994. Ribosomal DNA sequence analysis reveals new species grouping in the genus *Colletotricum*. *Experimental Mycology* 18: 121- 138.
- Sivan, A., Elad, Y. and Chet, I. 1984. Biological control effects of a new isolate of *Trichoderma harzianum* on *Pythium aphanidermatum*. *Phytopathology* 74: 498-501.
- Tianhui, Z. and Dexun, Q. 1994. Antagonism of *Trichoderma harzianum* to *Rhizoctonia solani*. *Journal Sichuan Agricultural University*. 12: 11-11.
- Van Belkum, A., Quint, W.G., de Pauw, B.E., Melchers, W.J.G. and Meis, J.F. 1993. Typing of *Aspergillus* species and *Aspergillus fumigatus* isolates by interrepeat polymerase chain reaction. *Journal Clinical Microbiology*. 31: 2502-2505.
- Ward, J.H. 1963. Hierarchical grouping to optimize an objective function. *Journal American Statistical Association*. 58: 236-244. doi: 10.2307/2282967.
- Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A. and Tingey, S.V. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* 18: 6531-6535.

*Submitted February 15, 2010– Accepted May 25, 2010*  
*Revised received June 11, 2010*